

$b\bar{b}b\bar{b}$ production in proton-nucleus collisions at the LHC

A. Del Fabbro and D. Treleani

*Dipartimento di Fisica Teorica dell'Università di Trieste and INFN, Sezione di Trieste,
Strada Costiera 11, Miramare-Grignano, I-34014 Trieste, Italy.*

Abstract

A sizable rate of events, with several pairs of b -quarks produced contemporarily by multiple parton interactions, may be expected at very high energies as a consequence of the large parton luminosities. The production rates are further enhanced in hadron-nucleus reactions, which may represent a convenient tool to study the phenomenon. We compare the different contributions to $b\bar{b}b\bar{b}$ production, due to single and double parton scatterings, in collisions of protons with nuclei at the CERN-LHC.

E-mail delfabbr@trieste.infn.it

E-mail daniel@trieste.infn.it

I. INTRODUCTION

The large rates of production of heavy quarks, expected at high energies, may lead to a sizable number of events, at the CERN-LHC, containing two or more pairs of b quarks, generated contemporarily by independent partonic collisions. An inclusive cross section of the order of $10 \mu\text{b}$ may in fact be foreseen for a double parton collision process, with two $b\bar{b}$ pairs produced in a pp interactions at 14TeV, while the cross section to produce two $b\bar{b}$ pairs by a single collision at the same c.m. energy may be one order of magnitude smaller [1]. All production rates are significantly enhanced in proton-nucleus collisions, which may offer considerable advantages for studying multiparton collision processes [2]. Given the large rates expected, the production of multiple pairs of b -quarks should hence represent a convenient process to study multiparton interactions in pA collisions at the LHC. On the other hand the mechanism of heavy quarks production is not yet understood satisfactorily also in the simplest case of nucleon-nucleon collisions, the effects of higher order corrections in α_s being still a matter of debate. A comprehensive description, of the much more structured process of heavy quarks production in hadron-nucleus interactions, might hence be approached after gaining a deeper understanding of the short scale parton-level dynamics of the process. On the other hand a significant feature of higher order corrections in α_s is that, for a limited set of physical observables, the whole effect of higher orders reduces to an approximate rescaling of a lowest order calculation of heavy quarks production in perturbative QCD [3]. Some of the features of the process are therefore effectively described by the simplest parton level dynamics at the lowest order in α_s , which lets one speculate that a similar property might hold also for a much more complex process, as $b\bar{b}b\bar{b}$ production.

Taking this optimistic point of view we will attempt, in the present note, to obtain indications on some properties of $b\bar{b}b\bar{b}$ production in pA interactions at the LHC, by considering the contributions of the two different interaction mechanisms, the connected $2 \rightarrow 4$ and the disconnected $(2 \rightarrow 2)^2$ parton processes, which will be effectively described by the lowest order diagrams in perturbative QCD, keeping higher order corrections into account by

a simple overall rescaling. Obviously, following this philosophy, all considerations will be limited to the class of physical observables falling in the category mentioned above.

II. $B\bar{B}B\bar{B}$ PRODUCTION IN PROTON-NUCLEUS COLLISIONS

Quite in general [4] [5], with the only assumption of factorization of the hard component of the interaction, the expression of the double parton scattering cross section to produce two $b\bar{b}$ pairs is given by

$$\sigma_{(p,A)}^D(b\bar{b}, b\bar{b}) = \frac{1}{2} \sum_{ij} \int \Gamma_p(x_i, x_j; s_{ij}) \hat{\sigma}(x_i, x'_i) \hat{\sigma}(x_j, x'_j) \Gamma_{(N,A)}(x'_i, x'_j; s_{ij}) dx_i dx'_i dx_j dx'_j d^2 s_{ij}, \quad (1)$$

where the index N refers to nucleon and A to a nucleus, while the indices i, j to the different kinds of partons that annihilate to produce a $b\bar{b}$ pair and the factor $1/2$ is a consequence of the symmetry of the expression for exchanging i and j . The interaction region of a hard process is very small as compared to the hadron scale, so the two interactions are well localized in transverse space within the two overlapping hadrons. The rate of events where two hard collisions take place contemporarily, in a given inelastic hadron-hadron interaction, depends therefore on the typical transverse distance between the partons of the pairs undergoing the multiple collision processes. A main reason of interest is hence that double parton scatterings may provide information on the typical transverse separation between pairs of partons in the hadron structure. Indeed the non-perturbative input of a double parton collision is the two-body parton distribution function $\Gamma(x_1, x_2, s_{1,2})$, which depends not only on the fractional momenta $x_{1,2}$, but also on the relative distance in transverse space $s_{1,2}$, besides (although not written explicitly to simplify the notation) on the scales of the two interactions and on the different kinds of partons involved. As a consequence the double parton scattering cross section is characterized by a linear dependence on dimensional scale factors, which have a direct relation with the typical transverse distances between the partons of the various pairs, contributing to the double scattering process under consideration.

The cross section is simplest when the target is a nucleon and partons are not correlated in fractional momenta, which may be a sensible approximation in the limit of small

x . In such a case the two-body parton distribution may be factorized as $\Gamma_p(x_i, x_j; s_{ij}) = G(x_i)G(x_j)F(s_{ij})$, where $G(x)$ are the usual one-body parton distributions and $F(s)$ a function normalized to 1 and representing the parton pair density in transverse space. The inclusive cross section to produce two pairs of b quarks is then written as [6]

$$\sigma_N^D(b\bar{b}; b\bar{b}) = \frac{1}{2} \sum_{ij} \Theta^{ij} \sigma_i(b\bar{b}) \sigma_j(b\bar{b}) \quad (2)$$

where $\sigma_i(b\bar{b})$ represents the inclusive cross sections to produce a $b\bar{b}$ pair in a hadronic collision, with the index i labelling a definite parton process. The factors Θ^{ij} have dimension an inverse cross section and result from integrating the products of the two-body parton distributions in transverse space. In this simplified case the factors Θ^{ij} provide a direct measure of the different average transverse distances between different pairs of partons in the hadron structure [6] [7].

The cross section has a more elaborate structure in the case of a nuclear target. The most suitable conditions to study the phenomenon are those where the nuclear distributions are additive in the nucleon parton distributions. In such a case one may express the nuclear parton pair density, $\Gamma_A(x'_i, x'_j; s_{ij})$, as the sum of two well defined contributions, where the two partons are originated by either one or by two different parent nucleons:

$$\Gamma_A(x'_i, x'_j; s_{ij}) = \Gamma_A(x'_i, x'_j; s_{ij})|_1 + \Gamma_A(x'_i, x'_j; s_{ij})|_2 \quad (3)$$

and correspondingly $\sigma_D^A = \sigma_D^A|_1 + \sigma_D^A|_2$. The two terms $\Gamma_A|_{1,2}$ are related to the nuclear nucleon's density. Introducing the transverse parton coordinates $B \pm \frac{s_{ij}}{2}$, where B is the impact parameter of the hadron-nucleus collision, one may write

$$\Gamma_A(x'_i, x'_j; s_{ij})|_{1,2} = \int d^2B \gamma_A(x'_i, x'_j; B + \frac{s_{ij}}{2}, B - \frac{s_{ij}}{2})|_{1,2} \quad (4)$$

where $\gamma_A|_{1,2}$ are given by

$$\begin{aligned} \gamma_A(x'_i, x'_j; B + \frac{s_{ij}}{2}, B - \frac{s_{ij}}{2})|_1 &= \Gamma_N(x'_i, x'_j; s_{ij})T(B) \\ \gamma_A(x'_i, x'_j; B + \frac{s_{ij}}{2}, B - \frac{s_{ij}}{2})|_2 &= G_N(x'_i)G_N(x'_j)T(B + \frac{s_{ij}}{2})T(B - \frac{s_{ij}}{2}) \end{aligned} \quad (5)$$

with $T(B)$ is the nuclear thickness function, normalized to the atomic mass number A and G_N nuclear parton distributions divided by the atomic mass number.

In the simplest additive case, the first term in Eq.(3) obviously gives a simple rescaling of the double parton distribution of an isolated nucleon

$$\Gamma_A(x'_i, x'_j; s_{ij})\Big|_1 = \Gamma_N(x'_i, x'_j; s_{ij}) \int d^2 B T(B) \quad (6)$$

and the resulting contribution to the double parton scattering cross section is the same as in a nucleon-nucleon interaction, apart from an enhancement factor due to the nuclear flux, which is given by the value of the atomic mass number A :

$$\sigma_A^D\Big|_1 = A\sigma_N^D. \quad (7)$$

The $\sigma_A^D\Big|_2$ term has more structure. In this case the integration on the relative transverse distance between the partons of the interacting pairs, s_{ij} , involves both the projectile and two different target nucleons:

$$\int ds_{ij} \Gamma_p(x_i, x_j; s_{ij}) T\left(B + \frac{s_{ij}}{2}\right) T\left(B - \frac{s_{ij}}{2}\right). \quad (8)$$

As one may notice in this case two very different scales are compared, the hadron radius r_p and the nuclear radius R_A . A usual approximation in pA interactions is to consider the limit $r_p \ll R_A$, where one may use the approximation

$$T\left(B \pm \frac{s_{ij}}{2}\right) \simeq T(B), \quad (9)$$

which allows one to decouple the integrations on s_{ij} and on B . One hence obtains:

$$\sigma_A^D\Big|_2 = \frac{1}{2} \sum_{ij} \int G_p(x_i, x_j) \hat{\sigma}(x_i, x'_i) \hat{\sigma}(x_j, x'_j) G_N(x'_i) G_N(x'_j) dx_i dx'_i dx_j dx'_j \int d^2 B T^2(B), \quad (10)$$

where

$$G_p(x_i, x_j) = \int d^2 s_{ij} \Gamma_p(x_i, x_j; s_{ij}). \quad (11)$$

Remarkably the two terms $\sigma_A^D\Big|_1$ and $\sigma_A^D\Big|_2$ have very different properties. In fact the correct dimensionality of $\sigma_A^D\Big|_1$ is provided by transverse scale factors related to the *nucleon*

scale, cfr. Eq.s(2),(7). The analogous dimensional factor in $\sigma_A^D|_2$ is provided by the *nuclear* thickness function, which is at the second power, being two the target nucleons involved in the interaction.

As pointed out in ref. [2], while on general grounds $\sigma_{(p,A)}^D$ depends both on the longitudinal and transverse parton correlations, the $\sigma_A^D|_2$ term depends solely on the longitudinal momentum fractions x_i, x_j so that, when the $\sigma_A^D|_2$ term is isolated, one has the capability of measuring the longitudinal and, a fortiori, also the transverse parton correlations of the hadron structure in a model independent way.

Although the two contributions may be defined also in a more general case, the separation of the cross section in the two terms $\sigma_A^D|_1$ and $\sigma_A^D|_2$ is most useful in the regime of additivity of the nuclear structure functions, which may not be a bad approximation for a sizable part of the kinematical regime of bottom quarks production at the LHC. In the case of a central calorimeter with the acceptance of the ALICE detector ($|\eta| < 0.9$), the average value of momentum fraction of the initial state partons, in a $pp \rightarrow b\bar{b}b\bar{b}$ process, is $\langle x \rangle \approx 6 \times 10^{-3}$. By introducing a cut in the transverse momenta of the b quarks of 5 GeV one obtains $\langle x \rangle \approx 10^{-2}$, while a cut of 20 GeV in p_t , within the same pseudorapidity range, gives $\langle x \rangle \approx 2 \times 10^{-2}$. If considering a more forward detector, as LHCb ($1.8 < \eta < 4.9$), the average value of momentum fraction is $\langle x \rangle \approx 5 \times 10^{-2}$. Deviations from additivity at low x are less than 10% for $x \geq 2 \times 10^{-2}$ [8] and, although increasing with the atomic mass number, non additive corrections are at most a 20% effect, on the kinematical regime above.

As already mentioned, heavy quarks production is characterized by a non trivial dynamics, in such a way that also next-to-leading corrections to the lowest order term in perturbative QCD are not sufficient for an exhaustive description of the inclusive spectra, which most likely need an infinite resummation to be evaluated. By comparing the results of different approaches to heavy quarks production (as NLO QCD and k_t -factorization) one nevertheless finds that, in a few cases, the whole effect of higher orders is to a large extent just an approximate rescaling of the results obtained by a lowest order evaluation in per-

turbative QCD. When limiting the discussion to an accordingly restricted set of physical observables, the whole effect of higher order corrections to heavy quark production is hence summarized by a single number, the value of the so-called K -factor. By evaluating, with the k_t -factorization approach, rapidity and pseudorapidity distributions of $b\bar{b}$ production, in pp collisions at the center-of-mass energy of $\sqrt{s} = 5.5$ TeV, within $|\eta| < 0.9$ and $1.8 < \eta < 4.9$, with different cuts in the transverse momenta of the produced b quarks, one finds a result not incompatible with a lowest order calculation in perturbative QCD rescaled by factor $K = 5.5$.

To discuss the production of $b\bar{b}b\bar{b}$ in pA collisions, while remaining in a kinematical regime where non additive corrections to the nuclear structure functions are not a major effect, we have limited all considerations to physical observables, where higher orders may be taken into account by the simple rescaling of the lowest order calculation. For $b\bar{b}b\bar{b}$ production, where only results of three level calculations are up to now available, we have further assumed that the value of the K -factor in the $2 \rightarrow 4$ parton process is the same as in the $2 \rightarrow 2$ process. We have hence evaluated the various contributions to the cross section in the case of a central calorimeter with the acceptance of the ALICE detector, $|\eta| < 0.9$, and of a more forward detector as LHCb, $1.8 < \eta < 4.9$. In both cases the typical values of x are rather small, so the factorization of the cross section in Eq.(2) may not be unreasonable. Since the process is dominated by gluon fusion, the expression of the cross section in Eq.(2) may be limited to a single term only. For the corresponding dimensional scale factor we have used the value reported by the CDF measurement of double parton collisions [9] [10]. To obtain the cross section at the lowest order in pQCD we have used the MRS99 parton distributions [11], with factorization and renormalization scale equal to the transverse mass of the b -quark. The cross sections of the $2 \rightarrow 4$ processes have been evaluated by generating the matrix elements of the partonic amplitudes with MadGraph [12] and HELAS [13]. For the mass of the bottom quark we have used the value $m_b = 4.6$ GeV. The multi-dimensional integrations have been performed by VEGAS [14] and all lowest order pQCD cross section have been finally multiplied by a factor $K = 5.5$.

III. RESULTS

A major result of the present analysis is that the effects induced by the presence of the nucleonic degrees of freedom, in double parton scatterings with a nuclear target, cannot be reduced to the simple shadowing corrections of the nuclear parton structure functions, whose effect is to decrease the cross section as a function of A . In the case of double parton collisions, the main effect of the nuclear structure is represented by the presence of the $\sigma_A^D|_2$ term in the cross section, which scales with a different power of A as compared to the single scattering contribution, producing an additive correction to the cross section.

To emphasize the resulting “anomalous” dependence of the double parton scattering cross section, as a function of the atomic mass number, we have plotted in fig.1 the ratio $\sigma_2^D/(\sigma_1^D + \sigma^S)$, as a function of A . The ratio represents the contribution to the cross section of the processes where two different nuclear target nucleons are involved in the interaction, scaled to the contribution where only a single target nucleon is involved. The dependence on the atomic mass number of the latter terms of the cross section, namely the single (σ^S) and the double (σ_1^D) parton scattering terms against a single nucleon in the nucleus, is the same of all hard processes usually considered, where nuclear effects may be wholly absorbed in the shadowing corrections to the nuclear structure functions. The contribution to the cross section of the σ_2^D term is, on the contrary, “anomalous”, involving two different target nucleons in the interaction process. The ratio above hence represents the relative weights of the “anomalous” to the “usual” contributions to the double parton scattering cross section on a nuclear target. The plots in fig.1 refer to the case of a forward calorimeter ($1.8 \leq \eta \leq 4.9$) and of a central calorimeter ($|\eta| \leq .9$) with different cuts on the transverse momenta of the produced b -quarks ($p_{cut} = 0, 5, 10$ GeV/c).

The different contributions to the cross section for $b\bar{b}b\bar{b}$ production, due to interactions with a single or with two different target nucleons, are shown in fig.2 in the case of a central calorimeter. The left figure shows the cross section as a function of the atomic mass number of the one-nucleon (dashed line) and of the two nucleons (continuous line) contributions

to the cross section. The dotted line is the sum of the two terms. The right figure shows the two contributions to the rapidity distribution of a b -quark produced in an event with two $b\bar{b}$ pairs: one-nucleon (dashed histograms) and two-nucleon contributions (continuous histograms) in the case of a heavy, Au , (higher histograms) and of a light nucleus, O , (lower histograms). The dependence on η of the two histograms is essentially the same, showing that the effect of the single parton scattering term (the $2 \rightarrow 4$ parton process) is negligible in this kinematical regime.

The A -dependence of the two different contributions, as a function of A , are shown in fig.3 in the case of a central calorimeter, after applying a cut of 5 GeV/c (left figure) and of 10 GeV/c (right figure) in the transverse momenta of each produced b -quark. Dashed, continuous and dotted lines have the same meaning as in fig.2.

The case of a forward calorimeter, $1.8 \leq \eta \leq 4.9$, is shown in fig.4, where the dashed, continuous and dotted lines have the same meaning as before.

Summarizing the large size of the cross section of $b\bar{b}b\bar{b}$ production in hadron-nucleus collisions at the LHC (the values are of the order of one hundreds of μb), suggests that the production of multiple pairs of b -quarks is fairly typical at high energies, hence representing a convenient channel to study multiple parton interactions. A rather direct feature, which is a simplest prediction and then a test of the interaction mechanism described above, is the “anomalous” dependence on A . The effects induced by the presence of the nucleonic degrees of freedom in the nuclear structure are in fact not limited, in this case, to the shadowing corrections to the nuclear structure functions usually considered, which cause a limited *decrease* (not larger than 20%, in the kinematical regime considered here) of the cross section for a hard interaction in hadron-nucleus collisions. When considering double parton scatterings, all nuclear effects can be exhausted in the shadowing corrections to the nuclear structure functions only in the $\sigma_A^D|_1$ term. The dominant effect of the nuclear structure is however due to the presence of the $\sigma_A^D|_2$ term in the cross section, which scales with a different power of A as compared to single scattering term, giving rise to a sizably larger correction, with opposite sign as compared to the shadowing correction, namely to an

increase of the cross section, which may become larger than 100% for a heavy nucleus.

Acknowledgment

This work was partially supported by the Italian Ministry of University and of Scientific and Technological Researches (MIUR) by the Grant COFIN2001.

FIGURES




FIG. 1. Relative weights of the terms with “anomalous” and “usual” A -dependence in the double scattering cross section for $b\bar{b}b\bar{b}$ production.




FIG. 2. Different contributions to the cross section for $b\bar{b}b\bar{b}$ production, in a central calorimeter. Left figure: cross section as a function of A of the one-nucleon (dashed line) and of the two nucleons (continuous line) contributions. The dotted line is the sum of the two terms. Right figure: differential rapidity distributions of a b -quark produced in an event with two $b\bar{b}$ pairs. One-nucleon (dashed histograms) and two-nucleon contributions (continuous histograms) in the case of a heavy (higher histograms) and of a light nucleus (lower histograms).




FIG. 3. Different contributions to the cross section for $b\bar{b}b\bar{b}$ production in a central calorimeter as a function of A , after applying a cut of 5 GeV (left figure) and of 10 GeV (right figure) in the transverse momenta of each produced b -quark: one-nucleon (dashed line), two nucleons (continuous line) processes and total (dotted line).




FIG. 4. Different contributions to the cross section for $b\bar{b}b\bar{b}$ production, in a forward calorimeter. Left figure: cross section as a function of A of the one-nucleon (dashed line) and of the two nucleons (continuous line) contributions. The dotted line is the sum of the two terms. Right figure: differential rapidity distributions of a b -quark produced in an event with two $b\bar{b}$ pairs. One-nucleon (dashed histograms) and two-nucleon contributions (continuous histograms) in the case of a heavy (higher histograms) and of a light nucleus (lower histograms).

REFERENCES

- [1] A. Del Fabbro and D. Treleani, Phys. Rev. **D66** (2002) 074012 [arXiv:hep-ph/0207311].
- [2] M. Strikman and D. Treleani, Phys. Rev. Lett. **88**, 031801 (2002) [arXiv:hep-ph/0111468].
- [3] M. G. Ryskin, A. G. Shuvaev and Y. M. Shabelski, Phys. Atom. Nucl. **64** (2001) 1995 [Yad. Fiz. **64** (2001) 2080] [arXiv:hep-ph/0007238].
- [4] N. Paver and D. Treleani, Nuovo Cim. A **70**, 215 (1982).
- [5] M. Braun and D. Treleani, Eur. Phys. J. C **18**, 511 (2001) [arXiv:hep-ph/0005078].
- [6] G. Calucci and D. Treleani, Phys. Rev. **D60** (1999) 054023.
- [7] A. Del Fabbro and D. Treleani, Phys. Rev. D **63**, 057901 (2001) [arXiv:hep-ph/0005273].
- [8] P. Amaudruz *et al.* [New Muon Collaboration], Nucl. Phys. B **441**, 3 (1995) [arXiv:hep-ph/9503291].
- [9] F. Abe *et al.* [CDF Collaboration], Phys. Rev. Lett. **79**, 584 (1997).
- [10] F. Abe *et al.* [CDF Collaboration], Phys. Rev. D **56**, 3811 (1997).
- [11] A.D.Martin, R.G.Roberts, W.J.Stirling and R.S.Thorne, Eur. Phys. J. **C14** (2000) 133.
- [12] T.Stelzer and W.F.Long, Comp.Phys.Comm. **81**, 357 (1994);
- [13] E.Murayama, I.Watanabe and K.Hagiwara, HELAS: HELicity Amplitude Subroutines for Feynman Diagram Evaluations, KEK report 91-11, January 1992;
- [14] G. P. Lepage, J. Comput. Phys. **27**, 192 (1978).